Grothendieck Polynomials of Inverse Fireworks Permutations

Jack Chou (UF) Tianyi Yu (UQAM)

March 18th

UCSD Combinatorics Seminar

Outline

1. Inverse Fireworks Permutations
2. Grothendieck Polynomials (Pipe Dreams)
3. Bumpless Vertical-less Pipe Dreams
4. Going-Up for Inverse Fireworks
5. Double Grothendieck Polynomials

Inverse Fireworks

Fireworks: The decreasing runs are increasing.

Inverse Fireworks

Fireworks: The decreasing runs are increasing.

3 1 6 7 5 4 2

Inverse Fireworks

Fireworks: The decreasing runs are increasing.

3 1 6 7 5 4 2

Inverse Fireworks

Fireworks: The decreasing runs are increasing.

3 1 6 7 5 4 2

3 < 6 < 7

Inverse Fireworks

Fireworks: The decreasing runs are increasing.

3 1 6 7 5 4 2

3 < 6 < 7



Inverse Fireworks

Fireworks: The decreasing runs are increasing.

3 1 6 7 5 4 2

3 < 6 < 7



3 1 7 5 6 4 2

Inverse Fireworks

Fireworks: The decreasing runs are increasing.

3 1 6 7 5 4 2



3 < 6 < 7

3 1 7 5 6 4 2

Inverse Fireworks

Fireworks: The decreasing runs are increasing.

3 1 6 7 5 4 2

3 < 6 < 7



3 1 7 5 6 4 2

3 < 7 ≮ 6

Inverse Fireworks

Fireworks: The decreasing runs are increasing.

3 1 6 7 5 4 2



3 < 6 < 7

3 1 7 5 6 4 2



3 < 7 ≮ 6

Inverse Fireworks

Why are they called

f*reworks?

Inverse Fireworks

Why are they called

f*reworks?







Inverse Fireworks

Fireworks ① 3__12 avoiding

Inverse Fireworks

Fireworks ① 3__12 avoiding

② Bell number $B(n)$

$3167542 \longleftrightarrow 1,3|2,7,4,5|6$

Inverse Fireworks

Fireworks ① 3__12 avoiding

② Bell number $B(n)$

$3167542 \longleftrightarrow 1,3|2,7,4,5|6$

Fireworks $\xleftrightarrow{\text{Inverse}}$ Inverse Fireworks

Grothendieck Polynomials and Pipe Dreams

Grothendieck Polynomials

K-theoretic analogues of Schubert polynomials

Grothendieck Polynomials

K-theoretic analogues of Schubert polynomials

$$G_w(x) = \begin{cases} x_1^{n-1} x_2^{n-2} \cdots x_{n-1} & \text{if } w = w_0 \\ \overline{\partial_i}\left(G_{ws_i}(x)\right) & \text{if } w(i) < w(i+1) \end{cases}$$

$$\partial_i(f) = \frac{f - s_i f}{x_i - x_{i+1}}$$

$$\overline{\partial_i}(f) = \partial_i\left((1 - x_{i+1}) f\right)$$

Grothendieck Polynomials

$$\partial_i(f) = \frac{f - s_i f}{x_i - x_{i+1}} \qquad \overline{\partial_i}(f) = \partial_i\left((1 - x_{i+1})f\right)$$

$$\overline{\partial_1}(x_1^2) =$$

Grothendieck Polynomials

$$\partial_i(f) = \frac{f - s_i f}{x_i - x_{i+1}} \qquad \overline{\partial_i}(f) = \partial_i\left((1 - x_{i+1})f\right)$$

$$\overline{\partial_1}(x_1^2) = \partial_1(x_1^2 - x_1^2 x_2) = \frac{(x_1^2 - x_1^2 x_2) - (x_2^2 - x_1 x_2^2)}{x_1 - x_2}$$


Grothendieck Polynomials

$$\partial_i(f) = \frac{f - s_i f}{x_i - x_{i+1}} \qquad \overline{\partial_i}(f) = \partial_i\left((1 - x_{i+1})f\right)$$

$$\overline{\partial_1}(x_1^2) = \partial_1(x_1^2 - x_1^2 x_2) = \frac{(x_1^2 - x_1^2 x_2) - (x_2^2 - x_1 x_2^2)}{x_1 - x_2}$$

$$= \frac{(x_1 + x_2)(x_1 - x_2) - x_1 x_2 (x_1 - x_2)}{x_1 - x_2} = x_1 + x_2 - x_1 x_2$$


Grothendieck Polynomials




Grothendieck Polynomials

Combinatorial formulas:

Pipe Dreams and Bumpless Pipe Dreams







Pipe Dreams










Pipe Dreams









Pipe Dreams







Pipe Dreams




Pipe Dreams




Pipe Dreams




Pipe Dreams




Pipe Dreams




Pipe Dreams




Pipe Dreams





Pipe Dreams



Pipe Dreams PD(2413) = set of pipe dreams of 2413









Pipe Dreams


PD(2413) = set of pipe dreams of 2413



$X_1 X_2^2$




$X_1^2 X_2^2$




$X_1^2 X_2$

Pipe Dreams


PD(2413) = set of pipe dreams of 2413



$X_1X_2^2$



$X_1^2X_2^2$



$X_1^2X_2$

$$\mathfrak{S}_{2413} = X_1^2X_2 + X_1X_2^2 - X_1^2X_2^2$$

Bumpless Vertical-less Pipe Dreams

Top Grothendieck

$\widehat{Gw}$ = highest degree terms of Gw

Top Grothendieck

$\widehat{G_w}$ = highest degree terms of G_w

1. Degree of $\widehat{G_w}$ determines the Castelnuovo-Mumford regularity of matrix Schubert varieties.
2. The support of $\widehat{G_w}$ conjecturally governs $\mathrm{Supp}(G_w)$.

Top Grothendieck

Theorem [PSW21]

$\forall w \in S_n$. $\exists!$ inverse fireworks permutation u such that

$$\widehat{G_w} = \lambda_w \cdot \widehat{G_u}$$

Top Grothendieck

Theorem [PSW21]


$\forall w \in S_n$. $\exists!$ inverse fireworks permutation u such that

$$\widehat{G_w} = \lambda_w \cdot \widehat{G_u}$$


$$\lambda_w = \widehat{G_v}(1, 1, \cdots, 1)$$

v inverse fireworks


BVPD




BVPD



BVPD





BVPD



Bumpless Vertical-less Pipe Dream


BVPD







BVPD







BVPD





BVPD





$w = 164235$

BVPD

Theorem [CY24]

For w inverse fireworks permutations,


$$\hat{G}_w = \sum_{D \in \mathrm{BVPD}(w)} X^{\mathrm{wt}(D)}$$

BVPD

Theorem [CY24]:


For w inverse fireworks permutations,

$$\hat{G}_w = \sum_{D \in \mathrm{BVPD}(w)} X^{\mathrm{wt}(D)}$$


$\mathrm{wt}(D)$ = # of  in each row

BVPD

$w = 164235$ $w^{-1} = 145362$




BVPD




BVPD


$x_1^4 x_2^2 x_3^2$




$x_1^4 x_2^3 x_3^1$




$x_1^3 x_2^4 x_3^1$



$x_1^3 x_2^3 x_3^2$







$x_1^2 x_2^4 x_3^2$

BVPD


$x_1^4 x_2^2 x_3^2$




$x_1^4 x_2^3 x_3^1$




$x_1^3 x_2^4 x_3^1$



$x_1^3 x_2^3 x_3^2$







$x_1^2 x_2^4 x_3^2$

$$\widehat{G_W}(X) = x_1^4 x_2^2 x_3^2 + x_1^4 x_2^3 x_3^1 + x_1^3 x_2^4 x_3^1 + x_1^3 x_2^3 x_3^2 + x_1^2 x_2^4 x_3^2$$


BVPD




BVPD




BVPD




BVPD




BVPD




BVPD




BVPD




BVPD




BVPD




BVPD




BVPD




BVPD



BVPD





BVPD



BVPD

PD is heighest weight if and only if

1) Blue pipes have no  or 

2) Blue pipes in first column are 

Going-Up for Inverse Fireworks

Support of G_w

$$\text{Supp}(G_w) = \{\text{monomials in } G_w\}$$ ~~coefficients~~

Support of G_w

$$\mathrm{Supp}(G_w) = \{\text{monomials in } G_w\}$$ ~~coefficients~~

Big Important Conjecture [HMMSD 22]:

Homogenized Grothendieck polynomials

are Lorentzian.

Support of G_w

$$\mathrm{Supp}(G_w) = \{\text{monomials in } G_w\}$$ ~~coefficients~~

Big Important Conjecture [HMMSD 22]:

~~Homogenized~~ Grothendieck polynomials

are Lorentzian.

Support of G_w

$$\text{Supp}(G_w) = \{\text{monomials in } G_w\}$$ ~~coefficients~~

Big Important Conjecture [HMMSD 22]:

~~Homogenized~~ Grothendieck polynomials

are ~~Lorentzian~~. "nice"

Support of G_w Conjectures [MSSD24]

1. If $m \in \mathrm{Supp}(G_w)$ not highest degree, then $\exists\, m' \in \mathrm{Supp}(G_w)$ of higher degree s.t. $m \mid m'$.

Support of G_w Conjectures [MSSD 24]

1. If $m \in \mathrm{Supp}(G_w)$ not highest degree, then $\exists\, m' \in \mathrm{Supp}(G_w)$ of higher degree s.t. $m \mid m'$.

2. If $m \in \mathrm{Supp}(G_w)$ not highest degree, then $\exists\, x_i$ such that $x_i m \in \mathrm{Supp}(G_w)$.

Support of G_w Conjectures [MSSD 24]

1. If $m \in \mathrm{Supp}(G_w)$ not highest degree, then $\exists\, m' \in \mathrm{Supp}(G_w)$ of higher degree s.t. $m \mid m'$.

2. If $m \in \mathrm{Supp}(G_w)$ not highest degree, then $\exists\, x_i$ such that $x_i m \in \mathrm{Supp}(G_w)$. (Going Up Conjecture)

Support of G_w Conjectures [MSSD 24]

1. If $m \in \mathrm{Supp}(G_w)$ not highest degree, then $\exists\, m' \in \mathrm{Supp}(G_w)$ of higher degree s.t. $m \mid m'$.

2. If $m \in \mathrm{Supp}(G_w)$ not highest degree, then $\exists\, x_i$ such that $x_i m \in \mathrm{Supp}(G_w)$. (Going Up Conjecture)

3. If $m_1, m_2 \in \mathrm{Supp}(G_w)$ and $m_1 \mid m_3 \mid m_2$, then $m_2 \in \mathrm{Supp}(G_w)$

Support of G_w Conjectures [MSSD 24]

1. If $m \in \mathrm{Supp}(G_w)$ not highest degree, then $\exists\, m' \in \mathrm{Supp}(G_w)$ of higher degree s.t. $m \mid m'$.

2. If $m \in \mathrm{Supp}(G_w)$ not highest degree, then $\exists\, x_i$ such that $x_i m \in \mathrm{Supp}(G_w)$. (Going Up Conjecture)

3. If $m_1, m_2 \in \mathrm{Supp}(G_w)$ and $m_1 \mid m_3 \mid m_2$, then $m_2 \in \mathrm{Supp}(G_w)$


[Fireworks, vexillary, Zero-One, Inverse fireworks]

Going Up

Question:


Given a PD not of highest weight, how can we increase the weight of one of the rows by exactly one?

Going Up



Going Up

Droop moves




Going Up

Lemma:


If PD is not highest weight, then we must be able to change  or do a droop move.

After a droop move, either the weight doesn't change or one of the rows increases by 1.


Going Up




Going Up




Going Up




Going Up




Going Up




Going Up




Going Up





Going Up



Going Up







Going Up

If $m \in \text{Supp}(G_w)$ not highest degree, then $\exists\, x_i$ such that $x_i m \in \text{Supp}(G_w)$.

Going Up

If $m \in \mathrm{Supp}(G_w)$ not highest degree, then $\exists\, x_i$ such that $x_i m \in \mathrm{Supp}(G_w)$.

Thm [CY24]:

True for inverse fireworks permutations.



Double Grothendieck Polynomials

Double Grothendieck $G_w(X;Y)$


Double Grothendieck $G_w(x;y)$

$$G_w(x;y) = \sum_{D \in PD(w)} \prod_{\boxplus \in D} (x_i + y_j - x_i y_j)$$

Double Grothendieck $G_w(x;y)$

$$G_w(x;y) = \sum_{D \in PD(w)} \prod_{\boxplus \in D} (x_i + y_j - x_i y_j)$$


$$\hat{G}_w(x;y) = \sum_{D \in BVPD(w)} \prod_{(i,j) = wt} (x_i y_j)$$




Double Grothendieck $G_w(x;y)$


$$G_w(x;y) = \sum_{D \in PD(w)} \prod_{\square \in D} (x_i + y_j - x_i y_j)$$


$$\hat{G}_w(x;y) = \sum_{D \in BVPD(w)} \prod_{(i,j) = wt} (x_i y_j)$$





$$G_w(x;1) = G_w(x) \qquad \hat{G}_w(x;1) = \hat{G}_w(x)$$















$$\widehat{G_w}(1,y) = y_1 y_2 y_3 \, y_1 y_2 y_3 y_4 y_5 = y_1^2 y_2^2 y_3^2 y_4 y_5$$



$$\widehat{G_w}(1,y) = y_1 y_2 y_3 \, y_1 y_2 y_3 y_4 y_5 = y_1^2 y_2^2 y_3^2 y_4 y_5$$

$$\deg(G_w(x)) = \deg(\widehat{G_w}(x)) = \deg(\widehat{G_w}(1;y)) = 3+5 = 8$$

w Invfire $\Longrightarrow$ $\widehat{G_w}(1,Y)$ is a single monomial

w Invfire $\Longrightarrow$ $\widehat{G_w}(1, Y)$ is a single monomial

w fireworks $\Longrightarrow$ $\widehat{G_w}(X, 1)$ is a single monomial

w Invfire $\Longrightarrow$ $\widehat{G_w}(1,Y)$ is a single monomial

w fireworks $\Longrightarrow$ $\widehat{G_w}(X,1)$ is a single monomial

$\Longleftrightarrow$ $\widehat{G_w}(X) = G_u(X)$ Dominant

w Invfire $\Longrightarrow$ $\widehat{G_w}(1,Y)$ is a single monomial

w fireworks $\Longrightarrow$ $\widehat{G_w}(X,1)$ is a single monomial

$\Longleftrightarrow$ $\widehat{G_w}(X) = G_u(X)$ Dominant

$\Longleftrightarrow$ rajcode(w) is weakly decreasing

Thank You!



